

AD-A186 344

PROBABILISTIC APPROACH TO COMPUTATIONAL ALGORITHMS FOR  
FINDING STATIONARY... (U) FLORIDA STATE UNIV TALLAHASSEE  
DEPT OF STATISTICS M I TAKSAR ET AL. OCT 86

1/1


UNCLASSIFIED

FSU-STATISTICS-M751 AFOSR-TR-87-1844

F/G 12/3

NL





AD-A186 344

## REPORT DOCUMENTATION PAGE

1a. SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS													
2a. SECURITY CLASSIFICATION AUTHORITY NA		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited													
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA															
4. PERFORMING ORGANIZATION REPORT NUMBER(S) FSU Technical Report No. M751		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 87-1044													
6a. NAME OF PERFORMING ORGANIZATION Florida State University	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research/NM													
6c. ADDRESS (City, State and ZIP Code) Department of Statistics Tallahassee, FL 32306-3033		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448													
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620-85-C-0007													
8c. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC		10. SOURCE OF FUNDING NOS. <table border="1"><tr><td>PROGRAM ELEMENT NO. 6.1102F</td><td>PROJECT NO. 2304</td><td>TASK NO. A3</td><td>WORK UNIT NO.</td></tr></table>		PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304	TASK NO. A3	WORK UNIT NO.								
PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304	TASK NO. A3	WORK UNIT NO.												
11. TITLE (Include Security Classification) Probabilistic Approach to Computational Algorithms for Finding Stationary Distributions of Markov Chains															
12. PERSONAL AUTHOR(S) Michael Taksar and Winfried K. Grassmann															
13a. TYPE OF REPORT Technical Report	13b. TIME COVERED FROM TO	14. DATE OF REPORT (Yr., Mo., Day) October, 1986	15. PAGE COUNT 9												
16. SUPPLEMENTARY NOTATION															
COSATI CODES <table border="1"><tr><td>FIELD</td><td>GROUP</td><td>SUB. GR.</td></tr><tr><td></td><td></td><td></td></tr><tr><td></td><td></td><td></td></tr><tr><td></td><td></td><td></td></tr></table>		FIELD	GROUP	SUB. GR.										18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.													
19. ABSTRACT (Continue on reverse if necessary and identify by block number) <p>A number of important theorems arising in connection with Gaussian elimination are derived, using semi-regenerative analysis. The implications of these theorems to find steady-state solutions of Markov chains are analysed. The results obtained in this way are then applied to quasi birth-death processes.</p>															
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED													
22a. NAME OF RESPONSIBLE INDIVIDUAL F. Proschan/M. Hollander M. Crowley		22b. TELEPHONE NUMBER (Include Area Code) 767 4-3218 5023	22c. OFFICE SYMBOL AFOSR/NM												

**AFOSR-TM- 87 - 1044**

**PROBABILISTIC APPROACH TO COMPUTATIONAL ALGORITHMS FOR  
FINDING STATIONARY DISTRIBUTIONS OF MARKOV CHAINS**

by

**MICHAEL I. TAKSAR\*  
FLORIDA STATE UNIVERSITY  
TALLAHASSEE, FL 32306**

and

**WINFRIED K. GRASSMANN  
UNIVERSITY OF SASKATCHEWAN  
SASKATOON, SASKATCHEWAN**

**FSU TECHNICAL REPORT NO. M751  
AFOSR TECHNICAL REPORT NO. 86-203**

**OCTOBER, 1986**

**\*Research supported by the Air Force Office of Scientific Research under  
Grant Number F49620-85-C-0007.**

Probabilistic Approach to Computational Algorithms for Finding Stationary Distributions of Markov Chains.

Michael .I. Taksar and Winfried K. Grassmann

**Abstract**

A number of important theorems arising in connection with Gaussian elimination are derived, using semi-regenerative analysis. The implications of these theorems to find steady-state solutions of Markov chains are analysed. The results obtained in this way are then applied to quasi birth-death processes.

Introduction

Originally, the determination of stationary distributions in Markov chains was done, using completely algebraic arguments. Unfortunately, algebraic arguments often do not take advantage of the particular structure of stochastic matrices which form the coefficients of the equilibrium equations. Recently, a new trend emerged which looks into the probabilistic interpretation of the algorithms for solving equilibrium equations. The objective of this paper is to further this trend by deriving computational algorithms from purely probabilistic arguments. The arguments employed in this paper are based on the semi-regenerative structure of Markov chains. This approach gives better insights into formal manipulations of equilibrium equations and provides probabilistic interpretations of the coefficients obtained at each step of such manipulations. This allows one in turn to make connections and draw conclusions that would not be obvious otherwise. In particular, we derive a number of theorems leading to a better understanding of Gaussian elimination



A-1

a semi-regenerative processes. These theorems allow one to derive relationships between certain matrix-geometric solutions and Gaussian elimination Neuts [1981], Gaver et al. [1984].

From a Markov chain  $\{Y_n, n=0,1,\dots\}$  one can obtain a semi-regenerative process by recording the state of the chain only while it visits points of a given subset  $D$  of the state space of  $Y_n$ . Specifically, let  $T_n, n = 1,2,\dots$ , be the time of the  $n^{\text{th}}$  visit to  $D$  and let  $X_n$  be the position of the chain at  $T_n$ . The process  $\{(X_n, T_n), n=1,2,\dots\}$  is then a Markov renewal process (see Cinlar [1975] Chapter 10),  $\{X_n, n=1,2,\dots\}$  is a semi-regenerative process and the  $T_n$  are the semi-regenerative epochs of the Markov renewal process. Thus, if we consider

$$A_n = \{ Y_{T_{n-1}}, Y_{T_{n-1}+1}, \dots, Y_{T_n-1} \}$$

then the conditional distribution of the sequence  $(A_n, A_{n+1}, \dots)$ , given the past of the process up to  $T_{n-1}$ , depends only on  $X_{n-1}$ , and all  $A_n$  are conditionally independent, given  $(X, T)$ . The analysis of the behaviour of  $Y$ , from one semi-regenerative epoch to another produces the main relation between steady-state probabilities that is used for developing the algorithm. For simplicity, we assume that the Markov chain  $\{Y_n, n=0,1,\dots\}$  is irreducible and aperiodic, and that the state space of  $Y$  is  $E=\{0,1,2,\dots\}$ . Such a Markov chain reaches steady state (see Cinlar [1975], Chapter 8), a fact that is expressed by the following relation

$$Y_n \rightarrow Y.$$

Alternatively, one can write

$$P_i(Y_n = j) \rightarrow p_j, j = 0,1,\dots,$$

Where  $p_j$  is the distribution of  $Y$ , and  $P_i(\cdot) = P(\cdot | Y_0 = i)$ . The notation  $E_i$  must be understood in a similar way.

Let

$$D = \{0,1,2,\dots,d-1\}$$

and

$$T = \min\{m > 0: Y_m \in D\}. \quad (1)$$

Let  $1_i(Y_n)$  be the indicator function for  $Y_n=i$ , and let

$$v_{ij}^{(d)} = E_i\left\{\sum_{m=0}^T 1_j(Y_m)\right\} = E_i\#\{m: m < T \text{ and } Y_m = j\}. \quad (2)$$

The symbol  $\#$  denotes the cardinality of a set. The expression,  $\#\{m: m > T \text{ and } Y_m = j\}$ , in particular, is the number of times  $m$  meets the condition in question, and the quantity  $v_{ij}^{(d)}$  becomes the expected number of visits to state  $j$  prior to the exit from  $E-D$ , given one starts from the point  $i$ . The  $v_{ij}^{(d)}$  are related to the  $p_i$  according to the following theorem:

**Theorem 1:** Let  $(p_0, p_1, \dots)$  be the steady-state distribution of the Markov chain  $Y_n$ . Then

$$p_j = \sum_{i=1}^{d-1} v_{ij}^{(d)} p_i. \quad (3)$$

**Proof:** Let

$$\begin{aligned} T_1 &= T \\ T_{n+1} &= \min\{m > T_n: Y_m \in D\} \\ X_n &= Y_{T_n}. \end{aligned} \quad (4)$$

From these definitions, it follows that  $T_n$  is a stopping time, and that  $X_n$  can only assume the values  $0,1,\dots,d-1$ . As a consequence, the strong Markov property applies, which means that  $X_n$  is a Markov chain with the state space  $D = \{0,1,\dots,d-1\}$ . This chain is irreducible because the original chain  $Y_n$  is

irreducible. Let  $\nu_i$ ,  $i = 0, 1, \dots, d-1$  be the unique invariant distribution for  $Y_n$ .

Let  $m(i) = E_i\{T\}$  be the times between regeneration, and let

$$K_n(i, j) = P_i\{X_n = j, T > n\}$$

According to Çinlar [1975], Chapter 10, Theorem (6.12)

$$\lim_{n \rightarrow \infty} P_i\{Y_n = j\} = \sum_{k \in D} \nu_k \sum_{n=0}^{\infty} K_n(k, j) / \sum_{k \in D} m(k) \nu_k \quad (5)$$

We must mention Çinlar assumes that the times between the Markov-renewals have a continuous, aperiodic distribution. Since we are dealing with a discrete-time Markov renewal process, the distributions in question have in fact a periodicity of 1. However, the proof for the discrete case is basically identical to its continuous counterpart. The sum of the  $K_n(k, j)$  in (5) can be found as

$$\begin{aligned} \sum_{m=0}^{\infty} K_m(i, j) &= \sum_{m=0}^{\infty} P_i\{Y_m = j, T > m\} \\ &= \sum_{m=0}^{\infty} E_i(1_j(Y_m) 1_{T > m}) = \nu_{ij}^{(d)}. \end{aligned} \quad (6)$$

Since  $P_i\{Y_n = j\}$  converges to  $p_j$ , one finds from (5) and (6)

$$p_j = \sum_{i=0}^d \nu_{ij}^{(d)} \nu_i / \sum_{k=0}^d m(k) \nu_k. \quad (7)$$

If  $i, j \in D$ ,  $\nu_{ij}^{(d)}$  is 1 if  $i=j$ , and zero otherwise. In this case, (6) becomes

$$p_i = \nu_i / \sum_{k=0}^d m(k) \nu_k, \quad (8)$$

a relation that is also well known from semi-Markov processes. From (7) and (8), one easily obtains (3), which proves the theorem. (3) can also be proven in different ways (Miller [1984], Grassmann et al. [1985]), but then, the connec-



tion to semi-regenerative theory is lost.

Later on, we need to relate the  $v_{ij}^{(k)}$  for different values of  $i, j, k$ . Such relationships can be obtained from the following theorem.

**Theorem 2:** Let  $v_{ij}^{(k)}$  be given by (2), and let  $p_{ij}$  be the transition probabilities of the Markov chain  $Y_n$ . Then for  $k > d$

$$v_{ij}^{(d)} = v_{ij}^{(k)} + \sum_{\nu=d}^{k-1} v_{i\nu}^{(d)} v_{\nu j}^{(k)} \quad (9)$$

**Proof:** Let  $D$  consist of the first  $d$  points from 0 to  $d-1$ , and let  $T$  be defined by (1). Similarly, let  $K$  consist of the first  $k$  points from 0 to  $k-1$ , and let

$$\tau = \min\{m > 0: Y_m \in K\} \quad (10)$$

Since the first visit to  $K$  occurs prior to the first visit to  $D$ ,  $\tau < T$ , and one has

$$\begin{aligned} v_{ij}^{(d)} &= E_i(\# \{m > 0: Y_m = j, m < T\}) \\ &= E_i(\# \{m: Y_m = j, 0 < m < \tau\}) + E_i(\# \{m: Y_m = j, \tau < m < T\}) \\ &= v_{ij}^{(k)} + E_i(\# \{m: Y_m = j, \tau < m < T\}). \end{aligned} \quad (11)$$

We now consider the successive visits to  $K-D$ . Thus, let  $\xi(n)$  the  $n^{\text{th}}$  visit to  $K-D$ , that is,

$$\begin{aligned} \xi(1) &= \min\{m: m > 0, Y_m \in K-D\} \\ \xi(n) &= \min\{m: m > \xi(n-1), Y_m \in K-D\}. \end{aligned}$$

The  $\xi(n)$  are obviously stopping times, and  $\xi(n) \rightarrow \infty$  as  $n \rightarrow \infty$ . Moreover,

$$\tau = \min\{m > 0: Y_m \in K\} < \min\{m > 0: Y_m \in K-D\} = \xi(1). \quad (11)$$

In (10), the equality holds if the first visit to  $K$  is also a visit to  $K-D$ . In a similar way, one finds from (1) and (11), provided  $K \supset D$

$$\tau < T. \quad (12)$$

(12) holds as an equality if the first visit to  $D$  coincides with the first visit to  $K$ . In this case, the second term of (11) is zero. If, on the other hand,  $\tau < T$ ,

the first visit to K precedes the first visit to D, that is, the first visit to K is a visit to K-D. Hence,

$$\tau = \xi(1) < T,$$

and one finds under this condition

$$\begin{aligned} & E_i[ \sum_{m=\xi(1)}^{\tau-1} 1_j(Y_m) 1_{m < T} ] \\ &= E_i[ \sum_{\nu=1}^{\xi(\nu+1)-1} \sum_{m=\xi(\nu)}^{\tau-1} 1_j(Y_m) 1_{m < T} ] \\ &= E_i[ \sum_{\nu=1}^{\xi(\nu+1)-1} \sum_{m=\xi(\nu)}^{\tau-1} 1_j(Y_m) 1_{m < T} \sum_{n=d}^{k-1} 1_n(Y_{\xi(1)}) ] \\ &= E_i[ \sum_{n=d}^{k-1} [ \sum_{\nu=1}^{\xi(\nu+1)-1} \sum_{m=\xi(\nu)}^{\tau-1} 1_j(Y_m) 1_{m < T} 1_n(Y_{\xi(1)}) ] ] \quad (13) \end{aligned}$$

We now condition on  $Y_{\xi(\nu)} = u$ . Because of the strong Markov property for the stopping time on  $\xi(\nu)$ , one has

$$\begin{aligned} & E_i[ \sum_{m=\xi(\nu)}^{\xi(\nu+1)-1} 1_j(Y_m) 1_{m < T} 1_n(Y_{\xi(\nu)}) \mid Y_{\xi(\nu)} = u ] \\ &= E_i[ \sum_{m=0}^{\xi(1)-1} 1_j(Y_m) 1_{m < T} 1_n(Y_0) \mid Y_0 = u ] \\ &= E_n[ \sum_{m=0}^{\tau-1} 1_j(Y_m) 1_{n=u} ] = 1_{n=u} v_{nj}^{(k)} \quad (14) \end{aligned}$$

We now use (14) to write (13) as

$$\begin{aligned}
 & j-1 = \xi(\nu+1)-1 \\
 & E_1 \left[ \sum_{n=d}^{j-1} \sum_{\nu=1}^{\xi(\nu)-1} \sum_{m=\xi(\nu)}^{l_m(Y_{\xi(\nu)})} l_{\xi(\nu)}(Y_{\xi(\nu)}) v_{nj}^{(k)} \right] \\
 & k-1 = \\
 & E_1 \left[ \sum_{n=d}^{k-1} \sum_{m=0}^{T-1} l_m(Y_m) v_{nj}^{(k)} \right] = \\
 & \sum_{n=d}^{k-1} E_1 \left[ \sum_{m=0}^{T-1} v_{ni}^{(k)} l_n(Y_m) \right] \\
 & \sum_{n=d}^{\infty} v_{nj}^{(k)} v_{in}^{(d)}.
 \end{aligned}$$

In summary

$$E_1 \left[ \sum_{m=0}^{\infty} l_m(Y_m) v_{nj}^{(k)} \right] = \sum_{n=d}^{\infty} v_{in}^{(d)} v_{nj}^{(k)} \quad (15)$$

If we compare (15) with (11), we obtain (3).

Corollary: For  $i < d$

$$v_{ij}^{(d)} = v_{ij}^{(d+1)} + v_{id}^{(d)} v_{dj}^{(d+1)}. \quad (16)$$

A similar result was obtained in Grassmann et al [1983].

### Applications

The theorems above are particularly useful if the states can be partitioned into groups of states  $K_0, K_1, K_2, \dots$ , called levels. Level 0 consists of state 0 only, and level  $n$ ,  $n=0,1,2,\dots$  consists of the states  $k_{n-1}$  to  $k_n-1$ . For consistency,  $k_0$  is defined as 0. If  $d$  in equation (3) is replaced by  $k_d$ , one has

$$p_j = \sum_{i=0}^{k_d-1} p_i v_{ij}^{(k_d)} = \sum_{n=0}^{d-1} \sum_{i \in K_n} p_i v_{ij}^{(k_d)} \quad j \in K_d \quad (17)$$

We now define the following vectors and matrices:

$$q_n = [ p_j, j \in K_n ]$$

and

$$U_{nd} = [ v_{ij}^{(k_d)}, i \in K_n, j \in K_d ] \quad (18)$$

Using these symbols, (17) becomes

$$q_d = \sum_{n=0}^{d-1} q_n U_{nd} \quad (19)$$

In many applications, one has the following: For all  $d$  within a certain range, say  $d > c$ , level  $d$  cannot be reached from level  $n$ ,  $n < d-1$ . In other words, level  $d-1$  cannot be skipped. Such levels will be called non-skippable. If  $d$  is

non-skippable,  $U_{d-2}, U_{d-3}, \dots$  are all zero, because all  $v_{ij}^{(k_d)}$  vanish. In this case, (19) becomes

$$q_d = q_{d-1} U_{d-1,d} \quad (20)$$

Suppose now that for  $n$  within the range extending from a certain lower limit  $c$  up to  $\infty$ , all levels have the following properties:

- 1) Level  $d$  is non-skippable
- 2)  $k_{d+1} = k_d + a$ , where  $a$  is a constant
- 3)  $[p_i]_{j+i, i \in K_d} = [p_i]_{j+i, i \in K_g}, j > 0, g > d$ .

It is easy to verify that under these conditions,  $U_{d-1,d} = U_{g-1,g} = U, g > d$ . From (20) one concludes that under these conditions,  $q_d$  is matrix-geometric (Wallace [1969], Neuts [1981], Gaver et al. [1984]), that is

$$q_{d+r} = q_d U^r. \quad (21)$$

Thus, using the  $v_{ij}^{(d)}$  allows one to derive a number of results. Moreover, the  $v_{ij}^{(d)}$  can be found in many cases of practical importance. Grassmann et al [1985] showed how to find the  $v_{ij}^{(j)}$ , using a modification of Gaussian elimination. Once the  $v_{ij}^{(j)}$  are determined, one can obtain the  $v_{ij}^{(d)}$ ,  $d < j$ , either using (9) or (16). Moreover, the  $v_{ij}^{(d)}$  can be grouped into matrices according to equation (18), which allows one to set up interesting relations for quasi birth-death processes, in particular equations (20) and (21).

#### References

- E. Çinlar (1975), Introduction to Stochastic Processes, Prentice-Hall Inc.
- D.P. Gaver, P.A. Jacobs, G. Latouche, Finite Birth-and-Death Models in Randomly Changing Environments, Adv. Appl. Prob. 16, pp 715-731.
- W.K. Grassmann, M.I. Taksar, D.P. Heyman (1985), Regenerative analysis and Steady State Distributions for Markov Chains, Operations Research, vol 33, pp 1107-1116.
- D.R. Miller (1981), Computation of Steady-State Probabilities for M/M/1 Priority Queues, Operations Research 29, pp 945-958.
- M.F. Neuts (1981), Matrix-Geometric Solutions in Stochastic Models, The John Hopkins University Press, Baltimore.
- V. Wallace (1969), The Solution of Quasi Birth and Death Processes Arising from Multiple Access Computer Systems, Ph.D. Dissertation, University of Michigan.

END

12-87

DTIC